

Some Causes of Genotypic and Phenotypic Discordance in Monozygotic Twin Pairs

Geoffrey A. Machin

Department of Laboratory Medicine and Pathology, University of Alberta Hospitals, Edmonton, Alberta, Canada

The use of the adjective “identical” rather than monozygotic leads to misunderstandings about the biology of monozygotic twinning. Most monozygotic twin pairs are not identical; there may be major discordance for birth weight, genetic disease, and congenital anomalies. These indicate that post-zygotic events may lead to the formation of two or more cell clones in the inner cell mass and early embryo that actually stimulate the monozygotic twinning event. There is also evidence that there may be unequal allocation of numbers of cells to the monozygotic twins; this may have widespread implications for the cascade of developmental events during embryogenesis, formation, and vascularization of the placenta.

Large-scale zygosity testing at birth could be the template for analysis of twin outcomes and their biologic causes.

© 1996 Wiley-Liss, Inc.

KEY WORDS: diseases in twins, monozygotic twins, embryogenesis

INTRODUCTION

Monozygotic (MZ) twins continue to be analyzed for the prevalence and cause of multifactorial diseases. Such studies are based on the premises that MZ twin pairs are genetically “identical,” but were exposed to different postnatal environments. For instance, cohorts of MZ twin pairs are monitored for non-genetic influences on risks for developing diabetes and cardiovascular disease [Newman et al., 1990]. The assumption of such studies is that “. . . by comparing within genetically identical MZ twin pairs, they have eliminated genetic variability, thereby creating a pure culture of non-

genetic determinants—determinants they define as being ‘environmental exposure’ and ‘personal behaviors’ . . .” [Van Itallie and Stunkard, 1990]. However, the work of Bouchard et al. [1990] clearly indicates that “personal behaviors” of MZ twins reared apart are surprisingly similar. Attempts to dissect out genetic and environmental factors in skin nevus counts and their progression to melanoma are similarly flawed. There is strong correlation between nevus counts within MZ twin pairs, but not in dizygotic (DZ) twin pairs [Easton et al., 1992]. However, attempts to distinguish between genetic predisposition to nevi and the effects of sun exposure cannot be made, since a tendency to sun exposure might be genetically determined in MZ twins [Cantor, 1992].

While it is true that most of MZ twins are phenotypically very similar, there are significant numbers of MZ pairs who are neither genotypically nor phenotypically “identical.” Post-zygotic genetic events of various types and severity result in striking within-pair dissimilarity; indeed, it is proposed that, in some pairs, these post-zygotic events not only precede the twinning process, but may actually trigger it; if two recognizably different cell clones exist in one early zygote, the differences may be sufficient to cause mutual “recognition and repulsion,” with resulting MZ twinning. Such dissimilarities strike at the heart of twin studies, since they point to genetic dissimilarity in some, and perhaps many, MZ twin pairs.

Even if there are also extrinsic events that cause MZ twinning (such as traumatic rupture of the zona pellucida during hatching), an important characteristic is that the allocation of blastomeres to the two embryos may not be equal. Hence, the development of the twins may run non-parallel courses from the moment of the twinning event. Certain critical masses of cells may be necessary for the cascade of normal transcriptional events that leads to cell lineages and differentiation following gastrulation. It is easy to see how “vanishing twins,” fetus papyraceus, and holoacardius acephalus (twin reversed arterial perfusion) might result from such unequal allocations of cell populations. In the case of X-chromosome inactivation in female MZ twin pairs, there is good evidence for unequal allocation of blastomeres in some cases (see below).

It should also be noted that, for each MZ twin to approach DZ twin or singleton birth weight for a given

Received for publication March 25, 1994; revision received July 29, 1994.

Address reprint requests to Geoffrey A. Machin, M.D., Department of Laboratory Medicine and Pathology, 5B4.08 Mackenzie Health Sciences Centre, University of Alberta Hospitals, Edmonton, Alberta, Canada, T6G 2R7.

gestational age, the whole zygote needs to pass through one extra mitotic cycle. The implications for this on timing and differentiation are not clear. If there is unequal allocation of blastomeres, the smaller twin might need more than one extra mitotic cycle to achieve normal body weight.

A further point is that the formation of MZ twins must result in major reorientation of at least one of the major axes of the body plan. Given that higher vertebrates undergo axial orientation and regional specification largely by inductive (or regulative) cell-to-cell interactions whose effects begin after gastrulation, it seems likely that the first of the three axes (possibly dorso-ventral) [Bocklage, 1981, 1987] is laid down in the oocyte cytoplasm as a maternal effect, i.e., by regionally unequal topographic distribution of maternal mRNAs that specify the position of polar trophoblast; thereafter, the cranio-caudal axis of gastrulation is specified early, and the third axis, left-right, is probably specified inductively somewhat later, although the molecular parameters may be laid out before gastrulation. In MZ twinning, it is easy to see that cranio-caudal and dorso-ventral axes (or gradients) could be transmitted fairly faithfully from a single zygote to the two gastrulation processes; but left-right asymmetry might well be disturbed. This could account for some cases of abnormal heart looping, situs inversus/ambiguus/heterotaxy and proposed mirror-imaging in MZ twins.

By corollary, it would follow that all animals with a virtually autonomous (i.e., mosaic) regional specification of body regions would be unable to undergo MZ twinning, since this would result in reciprocal deficiencies in the body plans of the twins (e.g., one largely dorsal or animal, the other largely ventral or vegetal, etc.).

A further important factor is that two-thirds of MZ twins share a truly single, monochorionic placenta, with the potential for asymmetrical unequal sharing of venous return from the placental parenchyma. Thus, even the prenatal environment for monochorionic MZ twin pairs may be more dissimilar than that for dichorionic twins (see accompanying paper).

The purpose of this paper is to draw attention to the several known and hypothesized factors that may cause genotypic and phenotypic dissimilarity in MZ twin pairs. The facile use of MZ and DZ twins to distinguish "genetic" and "environmental" influences on development and disease is to be discouraged [Phillips, 1993]. Thorough documentation of zygosity and chorionicity by prenatal and perinatal diagnosis [Derom et al., 1991] allows certain parameters of similar and dissimilar development to be measured in MZ twins (*see accompanying paper* [Machin et al., 1995]). Careful study of MZ pairs who are dissimilar will yield important information on the likely causes and effects of MZ twinning.

SOME CLUES TO THE ETIOLOGY OF MZ TWINNING

The timing of the MZ twinning events has been linked to placentation. One-third of MZ twins have dichorionic placentas, separate or fused. This implies an early separation of two groups of blastomeres, with each twin subsequently developing its own trophoblast

and inner cell mass. Two-thirds of MZ twins are monochorionic; in these twins, the "decision" for twinning is made later and is limited to the inner cell mass after it has separated from the trophoblast. There is discordance between the number of inner cell masses and the number of trophoblasts. A small proportion of monochorionic MZ twins are also monoamniotic, indicating an even later twinning event, when the amnion has become distinct from the inner cell mass(es). Most conjoined twins are monochorionic monoamniotic. They have two body axes in one body, implying a very late and variably incomplete attempt at MZ twinning.

Thus, the four MZ types, identified by their placentation as being of different timing, could be caused by different events. The only major pointer is a steadily rising female excess from dichorionic to conjoined MZ twins (Fig. 1) [Derom et al., 1988; James, 1980; Gedda et al., 1981]. It seems likely that dichorionic MZ twins are fundamentally different from monochorionic MZ twins in their origin and biology, apart from the fact that they are spared the disadvantages of a monochorionic placenta. Monochorionic MZ twins are significantly at risk for non-genetic prenatal environmental influences that distinguish them from dichorionic MZ twins.

Potentially informative placental arrangements are to be found in MZ higher multiple gestations. Here there may be combinations of dichorionic and monochorionic placentations, indicating serial twinning




Fig. 1. Sex ratios in twins by zygosity, chorion, and amnion status. There is a steadily increasing female excess in MZ twins arising from later twinning events [based on data by Derom et al., 1988; James, 1980; Gedda et al., 1981].

events that have happened at different times after conception. Significant similarities and differences between monozygotic and dizygotic twins among MZ triplets or quadruplets could point to the timing and effects of the post-zygotic triggers of twinning (*see accompanying paper*) (Fig. 2) [Machin and Bamforth, 1996].

POST-ZYGOTIC GENETIC EVENTS

Two major types of post-zygotic genetic events are now known to distinguish some MZ twin pairs (although chorionicity is not always documented):

1. Chromosomal Mosaicism in MZ Twins

The situation is complicated by three factors, viz., the timing of the event (even into somatic regions of one twin), the placental status of the chromosomes (i.e., interacting effects of confined placental mosaicism and confined MZ twin mosaicism), and the presence or absence of interfetal anastomoses, which can superimpose blood chimerism, even among MZ twins.

MZ twins have been reported concordant for all four of the common human aneuploidies; 45,X [Reiss et al., 1993; Heydanus et al., 1993, case 22; Lin and Garver, 1988], 21-trisomy [Shapiro and Farnsworth, 1972; Rogers et al., 1982; Zellweger, 1968], 18-trisomy [Mulder et al., 1989; Schlessel et al., 1990], and 13-trisomy [Loevy et al., 1985]. In many of these pairs, there is discordance for the phenotypic expression of the aneuploidy, although mosaicism is not invoked. However, for 45,X, 21-trisomy and 13-trisomy, mosaicisms have been reported in MZ twin pairs to account for striking phenotypic discordance. Mosaicism is common among X-chromosome aneuploid singletons (some of whom may be sole survivors of MZ twin pairs). In MZ twin pairs discordant

for 45,X descent from 47,XXY, 47,XXX, 46,XY, and 46,XX zygotes has been reported (Table I). In many of these cases, fibroblast cell lines show complete dichotomy of the two cell lines between the embryos, and the presence of two cell lines in blood of one or both twins is presumed to be caused by MZ blood chimerism via vascular anastomoses in monozygotic placentas. Others have mosaicism in fixed tissues also. In some informative cases, non-disjunction events appear to have taken place at first post-zygotic division. Kurosawa et al. [1992] reported MZ twins with 47,XXY and 45,X karyotypes, without any evidence of a 46,XY cell line. Ross et al. [1969] reported on a pair of MZ twins with 47,XXX and 45,X cell lines without any residual 46,XX cells.

The following types of twin mosaicism could arise from a single 46,XY zygote by single post-zygotic non-disjunctional events: 46,XY and 45,X; 46,XY and 46,XX/45,X; 46,XY/45,X and 45,X; 46,XY/45,X and 46,XY/45,X; 47,XXY and 45,X; 47,XXY and 47,XXY/45,X; 47,XXY/45,X and 45,X; 47,XXY/45,X and 47,XXY/45,X; as well as mosaic combinations of the three cell lines, 47,XXY/46,XY/45,X. Similar considerations apply to 46,XX/45,X, 47,XXY/45,X, and 47,XXX/45,X mosaicism. In theory, two different non-disjunctional events could lead to MZ 46,XY and 46,XX embryos derived from a single 47,XXY zygote. X-chromosome non-disjunctions have been reported in MZ triplets [Dallapiccola et al., 1985; Landy et al., 1988]. The triplets reported by Dallapiccola et al. [1985] were MC, implying a relatively late post-zygotic non-disjunction.

45,X chromosome status can confer hemizygosity for X-linked disease, and symptomatic Duchenne muscular dystrophy may occur in singleton X-chromosome aneuploid mosaic patients [Bortolini et al., 1986]. Bonilla et al. [1990] reported on MZ twin sisters, one of whom had clinical and laboratory evidence of Duchenne muscular dystrophy. The asymptomatic sister had an affected son. 45,X/46,XX mosaicism was present in lymphocytes but not fibroblasts of the affected twin. The unaffected twin had 46,XX chromosomes in lymphocytes.

MZ twins discordant for trisomy 21 have been documented. Rogers et al. [1982] reported monozygotic male twins who were discordant for trisomy 21. Each was non-mosaic in fibroblast lines, but there was 46,XY/47,XY,1 21 chimerism in blood. Shapiro and Farnsworth [1972] described discordant females who were MZ by multiple blood group antigen testing. One twin was 46,XX and the other was a 46,XX/47,XX,1 21 mosaic (or chimera). Both twins had atrioventricularis communis.

One pair of monozygotic MZ male twins has been reported as discordant for trisomy-13 [Heydanus et al., 1993, case 2].

In all cases with an originally trisomic zygote, considerations of uniparental heterodisomy apply to the euploid twins.

MZ twin mosaicism has also been found for structural chromosome abnormalities. Fujimoto et al. [1991] described twins who were MZ by multiple antigen testing to a probability of 0.9977. The twins were both ab-




Fig. 2. MZ higher multiple gestations may show evidence of multiple twinning events occurring in time clusters or occurring successively over intervals of up to several days.

TABLE I. X-Chromosome Aneuploidy Mosaicism*

Author	Twin A fibroblasts	Twin B fibroblasts	Blood chimerism	MZ testing
Edwards et al., 1966	UT 45,X	NM 45,X	45,X/46,XY	
Karp et al., 1975	UT 45,X/46,XY	NM 2	45,X/46,XY	BGA
Schmidt et al., 1976	UT 45,X	NM 46,XY	None	BGA
Reindollar et al., 1987	UT 45,X/46,XY	NM 45,X/46,XY	?	MC
Arizawa et al., 1988	UT 45,X	NM 46,XY	None	
Turpin et al., 1961	UT 45,X	NM 46,XY	2	MC
Perlman et al., 1990	UT 45,X	NM 46,XY	?	MC
Wapner et al., 1990	UT 45,X	NM 46,XY	2	MC
Dallapiccola et al., 1985	UT 45,X	NM 3 2 (triplets) 46,XY		MC
Kurosawa et al., 1992	UT 45,X/47,XXY	45,X/47,XXY		RFLP
Deacon et al., 1980	TRAP, features UT 45,X	NF pump twin 46,XX	45,X/46,XX	MC
Pedersen et al., 1980	UT 45,X	NF 2		BGA-9932
Ross et al., 1969	UT 45,X/47,XXX	NF 45,X/47,XXX		BGA 70-99
Kaplowitz et al., 1991	UT 45,X	NF 46,XX	45,X/46,XX	RFLP
Neilson et al., 1982 (case 3)	UT 45,X	NF 46,XX		MC
Reiss et al., 1993	UT 45,X	NF 46,XX		RFLP

*NF, normal female phenotype; NM, normal male phenotype; UT, Ullrich-Turner phenotype; BGA, blood group antigens; RFLP, restrictive fragment length polymorphism.

normal, one having an Ullrich-Turner phenotype and a fibroblast chromosome constitution of 45,X. The co-twin was genitally male, and fibroblast chromosome analysis was 45,X (78%)/46,X,idi(Y)(p11) (22%). Both had blood chimerism. The structurally abnormal Y, present in the zygote, was presumably lost from a progenitor embryonic cell by anaphase lag. Watson et al. [1990] reported monochorionic MZ twins who were discordant for a Y/1 translocation with partial trisomy 1. Chromosomes from fibroblasts of the malformed twin were 46,XY/46,X,2 Y,1 der(Y), t(Y;1)(q12;q21). The co-twin had a 46,XY karyotype from lymphocytes. The mutational event was presumably post-zygotic. In a well-studied case, Lacassie et al. [1993] reported twins who were proven MZ by DNA mini-satellite analysis and who were discordant for a 13r chromosome. The normal twin was a 46,XX female by blood lymphocytes, as was the abnormal twin. However, the second twin had clear evidence of left lateralized somatic mosaicism, and skin fibroblasts from the left and right sides showed 13% and 2.6% of cells respectively with a 46,XX,2 13,1 13r karyotype. Tsukamoto et al. [1993] reported on MZ twins who were discordant for 7q2 syndrome. The phenotypically normal twin had a 46,XY karyotype, while the abnormal twin had a 46,XY,del(7)(q32→qter) chromosome constitution. There was lymphocyte chimerism. Juberg et al. [1981] reported monochorionic MZ twins discordant for 46,XY,del(10)(p11p15). It is notable that the placenta, although monochorionic, had two discs.

Several other cases of mosaicism for structural chromosome abnormalities in the context of twin reversed arterial perfusion are listed by Wolf et al. [1991].

The question of ploidy "mosaicism" (or chimerism) is complex. Both in singletons and twins, fertilization of a first polar body and of the ovum actually results in two zygotes, and hence, true chimerism, derived from two sperm but only one primary oocyte [Kennerknecht et al., 1991; Tuerlings et al., 1993; Bieber et al., 1981]. The twins reported by Bieber et al. [1981] were MC; a

normal 46,XY male and a holoacardiac 69,XXX fetus shared the same placenta, which had the necessary inter-fetal anastomoses to support the acardiac fetus. Using HLA haplotypes, it could be shown that the triploid acardiac was heterodisomic for the maternal contribution; this was derived from a first polar body with evidence of crossing over, since both maternal haplotypes were represented on the chromosome of the polar body. Hence, these twins, although monochorionic, were actually DZ, with true chimerism rather than mosaicism. Wulfsberg et al. [1991] reported monochorionic female twins who probably had twin transfusion syndrome. One twin was phenotypically normal, with 46,XX chromosomes. The other twin had 46,XX/69,XXX mosaicism (65%/35%) in skin fibroblasts with clear evidence of asymmetry for dysmorphism, including cutaneous pigimentary dysplasia.

2. Skewed X-Chromosome Inactivation in MZ Female Twins

Non-random X-chromosome inactivation of structurally normal X-chromosomes is one possible cause for expression of X-linked disease in females [Wadeluis et al., 1993]. Skewed X-chromosome inactivation has also been proposed as the cause of discordant phenotypic expression of X-linked diseases in female MZ twin pairs. Several reports of such discordance are now well documented (Table II). The hypothesis depends on the fact that there have been no reported cases of female MZ twins, obligatory or proven heterozygotes for X-linked disease, in whom either both or neither are affected. In all cases, there is phenotypic discordance. However, see also Goodship et al. [1996]. There is good evidence that the clinically affected twin has non-random inactivation predominantly of the X-chromosome carrying the wild gene, while the unaffected twin either has predominant inactivation of the X-chromosome carrying the mutant gene or has random X-inactivation. Some pairs with skewed/random X-inactivation showed clear evidence of unequal allocation of blastomeres to the twin embryos.

TABLE II. Non-Random X-Inactivation in Female MZ Twin Pairs With Clinical Expression of X-Linked Diseases*

Author	Disease	Inactivation detection method	Affected twin DNA source		Normal twin DNA source		Possible mechanism	Pedigree
			F	L	F	L		
Tuckerman et al., 1985	XLMR	BRdU incorporation	2	Sk to wild	2	Sk to mutant	RSk, pre-T	
Burn et al., 1986	DMD	Somatic cell hybrids	Sk to wild	2	Sk to mutant	2	RSk, pre-T	? de novo, gonadal mosaic
Richards et al., 1990	DMD	DXS255	Sk to wild	Sk to wild	Sk to mutant	Sk to mutant	RSk, pre-T	maternal carrier
Lupski et al., 1991	DMD	DXS255	2	Sk to wild	2	2	UBA, post-T	? de novo, normal twin has affected boy
Abbadi et al., 1992	DMD	DXS255	2	Sk to wild	2	Sk to mutant	RSk, pre-T	mother, sister carriers; affected brother
Jorgensen et al., 1992	RGCB	DXS255	Sk to wild	R	Sk to mutant	R	UBA post-T, blood chimerism	?
Winchester et al., 1992	Hunter disease	DXS255 (M27b)	Sk to wild	Sk to wild	2	R	UBA, post-T	Maternal carrier
Krayer et al., 1993	XLMR	DXS255 (M27b)	2	Sk to wild	2	Sk to mutant	RSk, pre-T	

*F, fibroblasts; L, leucocytes; XLMR, X-linked mental retardation; Sk, skewed; RSk, reciprocal skewed X-inactivation; pre-T, occurs before twinning; DMD, Duchenne muscular dystrophy; R, random; UBA, unequal blastomere allocation; post-T, occurs after twinning; RGCB, red-green color blindness. Placental chorionicity was not known in any of these cases.

In effect, there are two possible mechanisms (Fig. 3); in one, there is reciprocal skewed X-inactivation, which may result from the aggregation/segregation within the inner cell mass of two clones of cells with dissimilar X-inactivation status, followed by mutual repulsion from the oppositely inactivated clones. In the other mechanism, there may be random X-inactivation in the inner cell mass, but unequal allocation of blastomeres to the twins. In the twin who is allocated very few progenitor blastomeres, it is likely that, by chance, there will be skewed X-inactivation, while the co-twin, derived from a larger number of blastomeres, will show random or near random X-inactivation. In the first mechanism, X-inactivation would precede and (might trigger) MZ twinning; the second mechanism might be extrinsically imposed on the zygote after or independent of X-inactivation. This mechanism would result in unequal patch sizes in the twin pair [Nance, 1990; Zneimer et al., 1993]. Reported cases of non-random X-inactivation in female MZ twins are shown in Table II, according to method of DNA analysis and likely mechanism of distribution of X-inactivation.

An obvious question is whether the process of X-inactivation is normally skewed (reciprocally or otherwise) in MZ female twin pairs who are not heterozygous for a mutant X-linked genetic disease. If skewed X-inactivation is normally present, it could act as a trigger to the MZ twinning process, and account for the known excess of females among MZ twins, most marked among monochorionic MZ twin pairs. If unequal X-inactivation is not demonstrable in non-diseased female MZ twins, it would be necessary to suggest that the mutant gene itself is in some way the stimulus to non-random X-inactivation.

Another possibility is that apparently singleton females, with X-linked disease secondary to skewed X-inactivation [Wadelius et al., 1993], may actually be surviving MZ twins [Bundy, 1991]. The patterns and role of X chromosome inactivation in MZ twinning are further discussed by Trejo et al. [1994] and, in accompanying papers in this symposium, by Goodship et al. [1996] and Bamforth et al. [1996].

3. Post-Zygotic Dominant Gene Mutation

This author is not aware of any reports of proven post-zygotic mutations leading to discordance for autosomal or X-linked dominant single gene discordance in MZ twins. However, MZ twins may be discordant for severity of clinical expression of such diseases. Discordance for cloverleaf skull has been reported in three pairs of MZ twins with thanatophoric dysplasia [Horton et al., 1983; Serville et al., 1984; Young et al., 1989]. In a pair of MZ twins with neurofibromatosis (probably a de novo mutation), there was markedly different density of café-au-lait spots [Vaughn et al., 1981]. The twin with more numerous and larger café-au-lait spots developed an optic glioma at the age of two. The twins were of equal birth weight, but there is a suggestion here of discrepancy in the size of clones (or patches) of cells. In six pairs of affected MZ twins, Easton et al. [1993] found high concordance for density of café-au-lait spots and neurofibromata. Costa et al.




Fig. 3. Two possible mechanisms of non-random X-inactivation (XI). The maternal X-chromosome and outlines of cells with active maternal X chromosome are in bold. **a:** Random X-inactivation in the singleton zygote. **b:** XI does not occur immediately on syngamy; the result would be total non-random XI in the singleton and in the same or opposite directions in MZ twins; this degree of non-random XI has never been observed with structurally normal X chromosomes. **c:** XI might precede and trigger MZ twinning by a process of "cloning and repulsion"; the result would be reciprocal non-random XI, which has been observed in some cases (see Table III). **d,e:** Extrinsic splitting of the zygote after (and independent of) XI could result in random and non-random XI if the allocation of blastomeres were unequal. This could explain difference in patch sizes that have been observed in MZ female twins discordant for expression of X-linked genetic diseases.

[1990] reported MZ twins discordant for Aicardi syndrome. Since this is an X-linked dominant disease, non-random X-inactivation rather than mutation could explain the discordance.

MZ twins may be discordant for severity of expression of tuberous sclerosis. Gomez et al. [1982] reported on two pairs of male MZ twins with discordance for timing of onset and extent of multisystem involvement by tuberous sclerosis. A female pair of MZ twins [Kondo et al., 1991] showed, at age 20 years, marked differ-

ences in severity of tuberous sclerosis expression. Northrup et al. [1990] reported 7-year-old MZ male twins, one of whom had tuberous sclerosis while the other was phenotypically uninvolved. There is some indication of somatic mosaicism in the affected twin. In a pair of MZ twins described by Brilliant et al. [1990], high-resolution Southern blotting showed a single discordant hybridizing fragment. Again, this might be explained by a post-zygotic mutational event in one twin of this discordant pair.

Van Thienen and Van der Auwera [1994] reported a pair of twins who were proven MZ by DNA RFLPs and who were discordant for spondylocostal dysostosis, presumably on the basis of a post-zygotic mutation in one of the twins after separation of the two embryonic masses.

4. Post-Zygotic Recessive Gene Mutation

Walker [1950] reported cleft palate in one MZ twin and sporadic retinoblastoma in the co-twin. There was more marked fetal growth retardation in the twin with retinoblastoma.

5. Monozygotic Twins and Imprintable Genes

Fourteen pairs of MZ twins have been noted to be discordant for Wiedemann-Beckwith syndrome [Franceschini et al., 1993; Leonard and Johnson, 1993; Leonard et al., 1995, this volume]. The two pairs most recently reported were male, while the other twelve pairs were female. Two pairs of MZ twins have been found to be concordant for Wiedemann-Beckwith syndrome; one pair was female and the other of unspecified sex. In general, the affected twin has been heavier at birth, and there is incremental post-natal growth discordance. Although imprinting was invoked to explain the excess of discordantly affected female twins [Lubinsky and Hall, 1991], reports of discordant MZ male twin pairs reopen the question of etiology. Placentation is not known in all cases; of the two pairs reported by Leonard et al. [1996], one was monochorionic and one was dichorionic. It seems likely that the discordance might trigger MZ twinning. The relative risk of neoplastic complications in affected MZ twins is not known in comparison to singleton cases of Wiedemann-Beckwith syndrome.

MZ TWINS DISCORDANT FOR LATERAL ASYMMETRY (MIRROR-IMAGING)

This matter is confusing because there are clearly different degrees of asymmetry of viscera, varying from situs inversus totalis to right- and non-right-handedness. There does not appear to be an excess of situs inversus among MZ twins, nor of MZ twins among cases of situs inversus [Torgersen, 1950], although individual cases of discordance for situs inversus have been reported in MZ twins [Teeuw and Kok, 1992]. Handedness is complicated by the increased frequency of non-right handedness in twinning families [Bocklage, 1987]. Nevertheless, discordance for handedness is well known in MZ twins [Gedda et al., 1981]. In a detailed report, Lohr and Bracha [1992] describe MZ twins who were discordant for psychiatric illnesses. One twin was schizophrenic and the other had an affective disorder. CT scans showed mirror image differences in cerebral occipital lobe morphology. The schizophrenic twin was fully right handed while the co-twin was left handed. Dermatoglyphics were interpreted as showing greater similarity between the right hands of the twins and the left hands of the co-twins; however, this author finds the similarity of the right hand to right hand and left to left just as great. Meshkova [1992] could find no EEG evidence of mirror-imaging in MZ twins.

Viscero-atrial status, including isomerism (heterotaxy) and normal and abnormal cardiac looping, are important components of congenital heart disease; con-

genital heart disease is more prevalent in MZ twins than in DZ twins or singletons [Burn, 1991]. Based on small numbers, there is an association between right-sided occipital hair whorl, non-right-handedness, and heterotaxy (discordant MZ twin pairs) [Burn, 1991]. Conjoined twins offer the opportunity to consider disturbances of axis formation. In particular cephalothoracopagus (janiceps) and dicephalic conjoined twins have severely abnormal axis determination, with corresponding congenital heart disease [Machin, 1993; Machin et al., 1993; Seo et al., 1985; Giris et al., 1993].

As mentioned previously, abnormal re-establishment of left-right asymmetry might be expected to occur in MZ twins, but the situation remains unclear at present. Detailed studies are still required, including placentation as an index of timing.

DISCORDANCE FOR MAJOR MALFORMATIONS (ESPECIALLY OF THE MIDLINE) IN MZ TWINS

Congenital anomalies in MZ twins include conjoined twinning, twin reversed arterial perfusion, birth weight discordance secondary to monochorionic placentation, other vascular complications of monochorionic placentation, and mechanical deformation because of multiple adjacent fetuses [Fogel et al., 1965]. Putting these "special and limited" anomalies aside, there is evidence that true malformations (that also occur in singletons and DZ twins) are more prevalent in MZ twins.

The original paper of Schinzel et al. [1979] is compromised by its inclusion of many like-sexed twin pairs of unproven zygosity. Discordance for major malformation is rare even in the largest series of MZ twin pairs [Cameron et al., 1983; Myrianthopoulos, 1990]. However, there are some single reports in the literature to support the idea that discordance for neural tube defect, holoprosencephaly, tracheo-esophageal fistula, vertebral anomalies, anal atresia, fistula with esophageal atresia, radial and renal dysplasia (VATER) association, ventral body wall defects, cloacal dysgenesis, and symmelia do occur in probable and proven MZ twin pairs. It has to be said that concordance for major and multiple (especially midline) malformations is also reported in MZ twins. As with all MZ twin studies, it is difficult to reach valid conclusions about prevalence and chorionicity, since chorionicity and zygosity is more often assumed than proven. Thus, the literature remains unclear, and more rigorous criteria for proof of MZ status are needed in many case reports.

Of particular interest is the high prevalence of symmelia among like-sexed twins, some of whom are documented as MZ by monochorionic placentation or multiple antigen studies. However, many are of unknown zygosity. There appears to be a definite overlap between symmelia and twin reversed arterial perfusion, as documented by Stocker and Heifetz [1987] and reported, for instance, by Hendry and Kohler [1956]. It is interesting to note that monochorionic twin pairs who are discordant for symmelia and/or bilateral renal agenesis may be monoamniotic. In these pairs, the affected twin is spared the usual deformational consequences of oligohydramnios because the unaffected co-twin adequately fills the amniotic sac with fetal urine [Davies et al.,

1971; Heydanus et al., 1993; Kohler, 1972; Maurer et al., 1974; Betti and Traisman, 1971; Johnstone and Benirschke, 1976; Marras et al., 1983]. In the case reported by Maurer et al. [1974], one monoamniotic twin had bilateral renal agenesis without deformation, while the co-twin had unilateral renal agenesis with adequate fetal urine production.

The position is less clear for holoprosencephaly. In the case report of Burck et al. [1981], a twin pair were MZ by multiple antigen testing. Both had holoprosencephaly but of varying cerebral and facial severity. Twin A had definite lateral ventricles and persistent premaxillary anlage, whereas Twin B had alobar holoprosencephaly with median cleft lip and palate. Sperber and Machin [1987] reported monochorionic MZ fetuses dying of twin transfusion syndrome who were discordant for holoprosencephaly. Several other cases of like-sexed twin discordance for holoprosencephaly are reported, but MZ status is assumed rather than proven.

There are several reports of discordant MZ twins with oculoauriculovertebral dysplasia [Goldenhar complex], summarized by Boles et al. [1987]. Of 5 cases fulfilling criteria for MZ twinning, placentation was known to be monochorionic in 3 pairs and dichorionic in 2 pairs. Thus, vasculogenic mechanisms cannot satisfactorily explain all MZ cases. An early disturbance of cephalogenesis clearly occurs, and this may be provoked by the twinning event itself. Affected MZ twins are of lower birth weights than their co-twins [Boles et al., 1987].

It has been suggested that discordance in MZ twin pairs for amyoplasia [Hall et al., 1983] and aganglionosis [Siplovich et al., 1983] may be caused by microvascular events in early embryogenesis in the context of monochorionic placentation. However, all the general causes of maldevelopment (e.g., unequal allocation of blastomeres) could apply equally well.

Major abnormal blastogenesis may account for the frequently reported occurrence of discordance in MZ twins for lateral body wall defect and other forms of "amniotic band syndrome." Close apposition of the developing axes of the MZ twins could interfere with closure of the connections of intra- and extra-embryonic coelom. Illustrative cases were published by Fiedler and Phelan [1983], Herva and Karkinen-Jaaskelainen [1984], Kancherla et al. [1981], Khudr and Benirschke [1972], Worthen et al. [198], and Pysher [1980].

MZ twin studies of cerebral ventricular size show high concordance rates in comparison with DZ twins. However, in MZ twins discordant for schizophrenia, there is also discordant cerebral ventricular size within twin pairs [Reveley et al., 1982].

DISCORDANT FETAL GROWTH IN MZ MONOCHORIONIC TWINS—CONSIDERATIONS OF PLACENTAL VASCULAR ANATOMY

Little is known about mechanisms by which the autonomous fetal and placental circulations become linked. However, the monochorionic twin placenta is truly a single organ, to which at least two fetal circulations become linked in multiple monochorionic pregnancy. The two or more fetal umbilical arterial and venous systems may be connected to equal or unequal zones of placental perfusion, quite apart from any considerations of superficial placental vascular anastomoses. Thus, it is possible for various combinations of vascular perfusion zones to occur, as listed in Table III and diagrammed in Figure 4 [Machin et al., 1996].

In the absence of superficial arterio-arterial and/or veno-venous anastomoses, type 1 vascularity is likely to cause significant fetal growth discordance without twin-twin transfusion, type 2 will result in growth-concordant twins who are likely to go to term without complications, while types 3 and 4 are likely to result in prenatal twin-twin transfusion. This is because part of the arterial perfusion of the one twin (donor) returns to the larger, overlapping venous return zone of the other twin (recipient) [Bendon and Siddiqui, 1989]. The presence of arterio-arterial and veno-venous anastomoses further complicates these issues, and explains why the prenatal diagnostic criteria for twin-twin transfusion are uncertain [Danskin and Neilson, 1989; Wenstrom et al., 1992]. In general, arterio-arterial and veno-venous anastomoses act as direct inter-twin connections that can actually mitigate the effects of potential twin-twin transfusion (see accompanying paper).

In twin reversed arterial perfusion, there are large arterio-arterial and veno-venous anastomoses, and the acardiac twin receives perfusion almost entirely from the pump co-twin rather than from any part of the monochorionic placenta. Twin reversed arterial perfusion may result from major cardiac malformation in the acardiac twin, close insertions of the two cords with early

TABLE III. Some Patterns of Zonal Allocation of Arterial and Venous Perfusion in Monochorionic Twin Placentas

	Twin A	Twin B	Result
1	Larger arterial zone 1 larger venous zone	Smaller arterial zone 1 smaller venous zone	B growth retarded
2	Normal (equal) arterial zone 1 normal (equal) venous zone	Smaller arterial zone 1 normal or larger venous zone	Equal fetal growth, uncomplicated MC pregnancy
3	Larger arterial zone 1 normal or smaller venous zone	Smaller arterial zone 1 normal or larger venous zone	TTT, A donor, B recipient
4	Normal arterial zone 1 larger venous zone	Normal arterial zone 1 smaller venous zone	TTT, A recipient, B donor




Fig. 4. Arterial and venous zonal patterns in MC twin placentas and their implications for fetal growth and twin transfusion. **a:** Type 1. **b:** Type 2. **c:** Type 3. **d:** Type 4 as per Table IV.

“dominance” of the two umbilical circulations by the pump twin, or from failure of connection between the somatic and placental parenchymal circulations of the acardiac twin. The pump twin perfuses the acardiac fetus by retrograde flow in the umbilical vessels of the acardiac twin [Kirkinen et al., 1989]. A significant number of acardiac fetuses are chromosomally abnormal, with confined MZ twin mosaicism [reviewed by Wolf et al. [1991]. In these cases, cardiac malformation may cause cardiac failure and compensatory assumption of perfusional function by the pump twin. In the absence of superficial anastomoses, a vanishing fetus or fetus papyraceus would result. Symmelia and twin reversed arterial perfusion frequently co-exist [Stocker and Heifetz, 1987].

VANISHING TWINS AND FETUS PAPYRACEUS

There is an excess of monochorionic twin pairs among spontaneous abortuses [Livingstone and Poland, 1980; Uchida et al., 1983; Benirschke, 1961]. Intrauterine death of one twin fetus can occur at any gestational age in spontaneous multiple pregnancies. When fetal demise occurs early, often in the first trimester, a “vanishing twin” may result. There is no information on the proportion of vanishing twins that originated as MZ twins, although any of the conditions causing MZ twin discordance could cause vanishing twins.

Fetus papyraceus refers to compressed or flattened recognizable fetal remnants from fetal demise in the second trimester. The effect on the surviving co-twin

presumably depends largely but not entirely on zygosity and chorionicity. In the large series analyzed by Kindred [1944], 141 cases had information about chorionicity; 93 cases (66%) were DC, whereas 48 (34%) were monochorionic, predominantly monochorionic diamniotic. There is a particular association between fetus papyraceus and the presence in the monochorionic co-twin of aplasia cutis and bowel atresia [Wagner et al., 1990]. These anomalies are probably caused by acute hemodynamic changes in monochorionic vascular arrangements at the time of the death of one fetus. Nance [1981] has reported resolution of hydrops and hydramnios in the recipient of twin-twin transfusion following death and mummification of the donor twin.

Fetal death of one monochorionic twin in the third trimester (often secondary to twin-twin transfusion or twin reversed arterial perfusion) produces a different constellation of lesions in the survivor. Cerebral, renal, splenic, hepatic, pulmonary, and myocardial infarctions have been reported [Benirschke, 1961; Patten et al., 1989; Yoshida and Soma, 1986; Enbom, 1985]. Similar lesions may result from attempts at selective terminations of monochorionic pregnancies in which the circulation of the intended survivor is not adequately protected [Golbus et al., 1988; Donnenfeld et al., 1989].

PSEUDO-CONCORDANCE IN MZ TWINS

Reference has already been made to the concordant absence of oligohydramnios syndrome in MZ, mono-

chorionic, monoamniotic twins, only one of whom has fetal anuria [Kohler, 1972; Maurer et al., 1974; Betti and Traisman, 1971; Marras et al., 1983]. Discordance for congenital hypothyroidism in monochorionic twins was masked, at neonatal screening, by prenatal transfusion of thyroid hormones from the normal co-twin via placental anastomoses [de Zegher and Vanderschueren-Lodeweyckx, 1989]. In the case of monochorionic twins, fetal death or a-fetoprotein-producing anomaly of one fetus produces high amniotic a-fetoprotein levels in the sac of the co-twin [Stirrat et al., 1979; Winsor et al., 1987; Holbrook et al., 1987]; this is because a-fetoprotein can diffuse across a septum consisting of two amniotic layers only, but cannot do so when there is also intervening chorion, as in dichorionic twinning. Such "induced" high a-fetoprotein levels in the amniotic sacs of anatomically normal monochorionic co-twins have led to termination of both fetuses.

SUMMARY AND CRITIQUE

There is at present a destabilizing influence on twin studies. This is the recognition that there are many reasons why MZ twin pairs may be phenotypically and genotypically discordant. This paper has reviewed a number of prenatal environmental and genetic causes for such discordance. This realization throws in serious doubt any facile assumptions about "nurture and nature" in DZ and MZ twins; it clearly indicates that many MZ twins are not "identical"; but it also demonstrates that such discordance has much information to offer about mechanisms of MZ twinning and of early development of normal and abnormal singleton embryos.

While most MZ twins are remarkably similar, the unusual exceptions are intellectually stimulating and also a challenge to clinical management.

A MODEST PROPOSAL FOR MULTICENTRE TWIN STUDIES

As has been mentioned at several points in this paper, many classic publications about MZ twins and twinning are, in fact, lacking in the appropriate criteria for the diagnosis of MZ status. Like-sexed twins and the use of the Hardy-Weinberg law are no substitute for definitive diagnosis of MZ twinning.

In general, diagnosis of MZ twinning is simple. Monochorionic twins are MZ; like-sex twins can be tested for DNA restriction fragment length polymorphism differences, using placenta, cord, and membranes.

In order to discover the frequency and significance of discordant events in MZ twinning, it seems to this author that these minimal criteria for MZ status could be applied on a multi-centre basis for the organization of a multidisciplinary study of a large cohort of MZ twins. In this way, significant numbers of MZ twins, concordant and discordant for phenotypic and genotypic traits, could be followed from birth. Simple issues (such as the prognosis for growth catchup after birth weight discordance of various causes) could be studied in numbers not available from single centers.

The establishment of zygosity at birth not only answers the question most frequently asked by parents and relatives; it is also an issue of the "right to know,"

and has medical and educational consequences [Derom et al., 1991]. Among the medical consequences are the access to rejection-free transplanted organs, such as skin for burns [Westerveld et al., 1986], closure of neural tube defect [Barwick et al., 1993], and hypospadias repair [Donovan and Maizels, 1986], and parathyroid gland for post-thyroidectomy permanent hypoparathyroidism [Segerberg et al., 1992], renal transplantation [Tilney, 1986], testis transplantation [Silber, 1978], and bone marrow transplantation [Champlin et al., 1984]. There would be great benefits to genetic and, particularly, twin research if the simple matter of zygosity was adequately diagnosed at or before birth.

REFERENCES

- Abbadi N, Philippe C, Cherry M, Gilgenkrantz S, Tome F, Kaplan JC, Fardeau M (1992): Monozygotic twins discordant for Duchenne muscular dystrophy—evidence for mirror X-chromosome inactivation. *Proceedings: 7th International Congress on Twin Studies*, Tokyo, Japan.
- Arizawa M, Suehara N, Takemura T, Nakayama M (1988): Monozygotic twins discordant for sex. *Acta Obstet Gynecol Jpn* 40: 1479–1482.
- Bamforth F (1996): X-chromosome inactivation is mostly random in placental tissues of female monozygotic twins and triplets. *Am J Med Genet* 61:209–215.
- Barwick WJ, Oakes J, Goldberg JA (1993): Closure of myelomeningocele using skin from stillborn twin. *Ann Plast Surg* 30:171–174.
- Bendon RW, Siddiqui T (1989): Clinical Pathology Conference: Acute twin-to-twin transfusion. *Pediatr Pathol* 9:591–598.
- Benirschke K (1961): Twin placenta in perinatal mortality. *N Y State Med J* 61:1499–1508.
- Betti RT, Traisman HS (1971): Sirenomelia: A spectrum of related syndromes. *Clin Pediatr* 10:238–240.
- Bieber FR, Nance WE, Morton CC, Brown JA, Redwine FO (1981): Genetic studies of an acardiac monster: Evidence for polar body twinning in man. *Science* 213:775–777.
- Bocklage CE (1981): On the timing of monozygotic twinning events. In "Twin Research 3: Twin Biology and Multiple Pregnancy," Gedda L, Parisi P (eds.). New York: Alan R Liss, Inc. 3:155–165.
- Bocklage CE (1987): Invited Editorial Essay: Twinning, nonright-handedness, and fusion malformations: Evidence for heritable causal elements held in common. *Am J Med Genet* 28:67–84.
- Bocklage CE (1987): The organization of the oocyte and embryogenesis in twinning and fusion malformations. *Acta Genet Med Gemellol* 36:421–431.
- Boles DJ, Bodurtha J, Nance WE (1987): Goldenbar complex in discordant monozygotic twins: A case report and review of the literature. *Am J Med Genet* 28:103–109.
- Bonilla E, Younger DS, Change HW, Tantravahi U, Miranda AF, Medori R, DiMauro S, Warburton D, Rowland LP (1990): Partial dystrophic deficiency in monozygotic twin carriers of the Duchenne gene discordant for clinical myopathy. *J Neurol* 40:1267–1270.
- Bortolini ER, da Silva DM, Chequer RS, Vianna-Morgante AM, Katz M (1986): Duchenne muscular dystrophy in a girl with a 45,X/46,XX/47,XXX chromosome constitution. *Am J Med Genet* 25: 239–243.
- Bouchard TJ, Lykken DT, McGue M, Segal NL, Tellegen A (1990): Sources of human psychological differences. The Minnesota Study of Twins Reared Apart. *Science* 250:223–228.
- Brilliant MH, Cleve H, Edwards JH, Waldinger D (1990): Cell lines from monozygotic twins discordant for tuberous sclerosis display protein and DNA differences. *Am J Hum Genet* 47:A210.
- Bundey S (1991): Uses and limitations of twin studies. *J Neurol* 238:360–364.
- Burck V, Hayek HW, Zeidler V (1981): Holoprosencephaly in monozygotic twins—Clinical and computer tomographic findings. *Am J Med Genet* 9:13–17.
- Burn J, Povey S, Boyd Y, Munro EA, West L, Harper K, Thomas D (1986): Duchenne muscular dystrophy in one of monozygotic twin girls. *J Med Genet* 23:494–500.

- Burn J (1991): Disturbance of morphological laterality in humans. In "Biological Asymmetry and Handedness," Ciba Foundation Symposium 162. Bock GR, Marsh J (eds.). New York: John Wiley and Sons, pp 282-296.
- Cameron AH, Edwards JH, Derom R, Thierry M, Boelaert R (1983): The value of twin surveys in the study of malformations. *Eur J Obstet Gynecol Reprod Biol* 14:347-356.
- Cantor RM (1992): Analysis of nevus count data from twins. *Cytogenet Cell Genet* 59:185-187.
- Champlin RE, Feig SA, Sparkes RS, Gale RP (1984): Bone marrow transplantation from identical twins in the treatment of aplastic anemia: Implication for the pathogenesis of the disease. *Br J Haematol* 56:455-463.
- Costa T, Greer W, Duckworth-Rysiecki G, Musarella M, Ray P (1990): Monozygotic twins discordant for Aicardi syndrome. *Am J Hum Genet* 47[Suppl]:A53.
- Dallapiccola B, Stomeo C, Ferranti G, Di Lecce A, Purpura M (1985): Discordant sex in one of three monozygotic triplets. *J Med Genet* 22:6-11.
- Danskin FM, Neilson JP (1980): Twin-to-twin transfusion syndrome: What are the appropriate diagnostic criteria? *Am J Obstet Gynecol* 161:365-369.
- Davies J, Chazen E, Nance ME (1971): Symmelia in one of monozygotic twins. *Teratology* 4:367-378.
- Deacon JS, Machin GA, Martin JM, Nicholson S, Nwanko DC, Wintemute R (1980): Investigation of acephalus. *Am J Med Genet* 5:85-99.
- Derom C, Vlietinck R, Derom R, van den Berghe H, Thierry M (1988): Population-based study of sex proportion in monoamniotic twins. *N Engl J Med* 319:119-120.
- Derom C, Vlietinck R, Derom R, Keith LG, van den Berghe H (1991): Zygosity testing at birth: A plea to the obstetrician. *J Perinat Med* 19[Suppl]:234-240.
- de Zegher F, Vanderschueren-Lodeweyckx M (1989): Congenital hypothyroidism and monoamniotic twins. *Lancet* 2:169-170.
- Donnenfeld AE, Glazerman LR, Cutillo DM, Librizzi RJ, Weiner S (1989): Fetal exsanguination following intrauterine angiographic assessment and selective termination of a hydrocephalic, monozygotic co-twin. *Prenat Diagn* 9:301-308.
- Donovan J, Maizels M (1986): Transplantation of the prepuce to facilitate hypospadias repair in monozygotic twins. *J Urol* 136:1077-1079.
- Easton DF, Cox GM, MacDonald AM, Ponder BJ (1992): The study of nevi in British twins. Study design and description of the data set. *Cytogenet Cell Genet* 59:165-166.
- Easton DF, Ponder ME, Huson SM, Ponder BAJ (1993): An analysis of variation in expression of neurofibromatosis (NF) type I (NFI): Evidence for modifying genes. *Am J Hum Genet* 53:305-313.
- Edwards JH, Dent K, Kahn J (1966): Monozygotic twins of different sex. *J Med Genet* 3:117-123.
- Enbom JA (1985): Twin pregnancy with intrauterine death of one twin. *Am J Obstet Gynecol* 152:424-429.
- Fiedler JM, Phelan JP (1983): The amniotic band syndrome in monozygotic twins. *Am J Obstet Gynecol* 146:864-865.
- Fogel BJ, Nitowsky MM, Gruenwald P (1965): Discordant abnormalities in monozygotic twins. *J Pediatr* 66:64-72.
- Franceschini P, Guala A, Valden MP, Franceschini D (1993) (Letter): Monozygotic twins and Wiedemann-Beckwith syndrome. *Am J Med Genet* 46:353-354.
- Fujimoto A, Boelter WD, Sparkes RS, Lin MS, Battersby K (1991): Monozygotic twins of discordant sex both with 45,X/46,Xdic(Y) mosaicism. *Am J Med Genet* 41:239-245.
- Gedda L, Brenici G, Franceschini P, Talone C, Ziparo R (1981): A study of mirror imaging in twins. In "Twin Research 3: Twin Biology and Multiple Pregnancy." Gedda L, Parisi P, Nance WE (eds.). New York: Alan R Liss, pp 167-168.
- Gedda L, Nun MT, Carrega R, Roselli E (1981): Parental age, sibships' size, and sibships' sex ratio in twins: A progress report. In "Twin Research 3: Twin Biology and Multiple Pregnancy." New York: Alan R Liss, pp 75-80.
- Gerlis LM, Seo JW, Hos Y, Chi JG (1993): Morphology of the cardiovascular system in conjoined twins: Spatial and sequential segmental arrangements in 36 cases. *Teratology* 47:91-108.
- Golbus MS, Cunningham N, Goldberg JD, Anderson R, Filly R, Callen P (1988): Selective termination of multiple gestations. *Am J Med Genet* 31:339-348.
- Gomez MR, Kuntz NL, Westmoreland BF (1982): Tuberous sclerosis, early onset of seizures, and mental subnormality: Study of discordant monozygous twins. *Neurology* 32:604-611.
- Goodship J, Carter J, Burn J (1996): X-inactivation patterns in monozygotic and dizygotic female twins. *Am J Med Genet* 61:205-208.
- Hall JG, Reed SD, McGillivray BC, Herrmann J, Partington MW, Schinzel A, Shapiro J, Weaver DD (1983): Amyoplasia: Twinning in amyoplasia—A specific type of arthrogryposis with an apparent excess in discordantly affected identical twins. *Am J Med Genet* 15:591-599.
- Hendry DW, Kohler HG (1956): Sirenomelia ("mermaid"). *J Obstet Gynaecol Br Emp* 63:865-870.
- Herva R, Karkinen-Jasskelainen M (1984): Amniotic adhesion malformation syndrome: Fetal and placental pathology. *Teratology* 19:11-19.
- Heydanus R, Santema JG, Steward PA, Mulder PGH, Wladimiroff JW (1993): Preterm delivery rate and fetal outcome in structurally affected twin pregnancies: A retrospective matched control study. *Prenat Diagn* 13:155-162.
- Holbrook RH, Krovoza AM, Schelley S, Ferguson JE (1987): Biamniotic elevated alpha-fetoprotein and positive acetyl cholinesterase in twins, one with anencephaly. *Prenat Diagn* 7:653-655.
- Horton WA, Harris DJ, Collins DL (1983): Discordance for the Kleeblattschädel anomaly in monozygotic twins for thanatophoric dysplasia. *Am J Med Genet* 15:97-101.
- James WM (1980): Sex ratio and placentation in twins. *Ann Hum Biol* 7:273-276.
- Johnstone HB, Benirschke K (1976): Monozygotic twin discordant for urinary tract anomalies and presenting as hydramnios. *Obstet Gynecol* 47:610-615.
- Jorgensen AL, Philip J, Raskind WH, Matsushita M, Christensen B, Dreyer V, Motulsky AG (1992): Different pattern of X inactivation in MZ twins discordant for red-green color-vision deficiency. *Am J Hum Genet* 51:291-298.
- Juberg RC, Stallard R, Straughan WJ, Avotri KJ, Washington JW (1981): Clinicopathologic conference: A newborn monozygotic twin with abnormal facial appearance and respiratory failure. *Am J Med Genet* 10:193-200.
- Kancherla PL, Untawale VG, Gabriel JB, Chauhan PM (1981): Intrauterine amputation in one monozygotic twin associated with amniotic band: A case report. *Am J Obstet Gynecol* 140:347-348.
- Kaplowitz PB, Bodurtha J, Brown J, Spence JE (1991): Monozygotic twins discordant for Ullrich-Turner syndrome. *Am J Med Genet* 41:78-82.
- Karp L, Bryant JI, Tagatz G, Giblett E, Fialkow PJ (1975): The occurrence of gonadal dysgenesis in association with monozygotic twinning. *J Med Genet* 12:70-78.
- Kennerknecht J, Barbi G, Djalali M, Just W, Vogel W, Terinde R (1991): Uncommon chromosomal mosaicism in chorionic villi. *Prenat Diagn* 11:569-575.
- Khudr G, Benirschke K (1972): Discordant monozygous twins associated with amnion rupture: A case report. *Obstet Gynecol* 39:713-716.
- Kindred JE (1944): Twin pregnancies with one twin blighted: Report of two cases with comparative study of cases in the literature. *Am J Obstet Gynecol* 48:642-682.
- Kirkinen P, Herva R, Rasanen J, Airaksinen J, Ikaheimo M (1989): Documentation of paradoxical umbilical blood supply of an acardiac twin in the antepartum state. *J Perinat Med* 17:63-65.
- Kohler HG (1972): An unusual case of sirenomelia. *Teratology* 6:295-301.
- Kondo S, Yamashina U, Sato N, Asu K (1991): Discordant expression of tuberous sclerosis in monozygotic twins. *J Dermatol (Tokyo)* 18:178-180.
- Kruyer H, Mila M, Glover G, Castellini-Bel S, Carbonelli P (1993): Monozygotic twins discordant for the fragile X syndrome (FXS): *Am J Hum Genet* 53[Suppl 3]:A465.
- Kurosawa K, Kuromaru R, Imaizumi K, Nakamura Y, Ishikawa F, Ueda K, Kuroki Y (1992): Monozygotic twins with discordant sex. *Acta Genet Med Gemellol* 41:301-310.

- Lacassie Y, Cortes F, Molina V, Montgomery AH (1993): Discordant phenotype due to ring 13 chromosomal mosaic syndrome in one identical twin. *Am J Hum Genet* 53[Suppl 3]:A571.
- Landy HJ, Larsen JW, Schoen M, Larsen ME, Kent SG, Weingold AB (1988): Acardiac fetus in a triplet pregnancy. *Teratology* 37:1-6.
- Leonard NJ, Johnson CL (1993): Male monozygotic twins discordant for Beckwith-Wiedemann syndrome. *Am J Hum Genet* 53[Suppl 3]:A466.
- Leonard NJ, Bernier FP, Rudd N, Machin GA, Bamforth F, Bamforth S, Grundy P, Johnson C (1996): Two pairs of male monozygotic twins discordant for Beckwith-Wiedemann syndrome. *Am J Med Genet* 61:253-257.
- Lin AE, Garver KI (1988): Monozygotic Turner syndrome twins: Correlation of phenotypic severity and heart defect. *Am J Med Genet* 29:529-531.
- Livingstone JE, Poland BJ (1980): A study of spontaneously aborted twins. *Teratology* 21:139-148.
- Loevy HT, Miller M, Rosenthal IM (1985): Discordant monozygotic twins with trisomy 13. *Acta Genet Med Gemellol* 34:185-188.
- Lohr JB, Bracha HS (1992): A monozygotic mirror-image twin pair with discordant psychiatric illnesses: A neuropsychiatric and neuro-developmental evaluation. *Am J Psychiatry* 149:1091-1095.
- Lubinsky MS, Hall JG (1991): Genomic imprinting. Monozygous twinning and X-inactivation. *Lancet* 337:1288.
- Lupski JR, Garcia GA, Zoghbi HY, Hoffman EP, Fenwick RG (1991): Discordance of muscular dystrophy in monozygotic female twins: Evidence supporting asymmetric splitting of the inner cell mass in a manifesting carrier of Duchenne dystrophy. *Am J Med Genet* 40:354-364.
- Machin GA, Sperber GM, Ongaro I, Dabbagh J, Murdoch CA (1993): Cephalothoracopagus (Janiceps) conjoined twins: Computerized three-dimensional reconstruction. In *Blastogenesis Normal and Abnormal*. Opitz JM (ed). March of Dimes Birth Defects Foundation. Birth Defects: Original Article Series 29(1):253-259.
- Machin GA (1993): Conjoined twins: Implications for blastogenesis. In *Blastogenesis Normal and Abnormal*. Opitz JM (ed). March of Dimes Birth Defects Foundation. Birth Defects: Original Article Series 29(1):141-179.
- Machin GA, Bamforth F, Innes M, McNicol K (1995): Some perinatal characteristics of monozygotic twins who are dichorionic. *Am J Med Genet* 55:71-76.
- Machin GA, Bamforth F (1996): Zygosity and placental anatomy in 15 consecutive sets of spontaneously conceived triplets. *Am J Med Genet* 61:247-252.
- Machin GA, Still K, Lalani T (1996): Correlations of placental vascular anatomy and clinical outcomes in 69 monochorionic twin pregnancies. *Am J Med Genet* 61:229-236.
- Marras A, Mereu G, Dessi C, Macciotta A (1983): Oligohydramnios and extrarenal abnormalities in Potter syndrome. *J Pediatr* 102:597-598.
- Maurer SM, Dobrin RS, Vernier RL (1974): Unilateral and bilateral agenesis in monoamniotic twins. *J Pediatr* 84:236-238.
- Meshkova TA (1992): Laterality effects in twins. *Acta Genet Med Gemellol* 41:325-333.
- Mulder AFP, van Eyck J, Groenendaal F, Wladimiroff JW (1989): Trisomy 18 in monozygotic twins. *Hum Genet* 83:300-301.
- Myrianthopoulos NC (1970): An epidemiologic survey of twins in a large, prospectively studied population. *Am J Hum Genet* 22:611-629.
- Nance WE (1981): Malformations unique to the twinning process. In Gedda L, Parisi P, Nance WE (eds): "Twin Research 3: Twin Biology and Multiple Pregnancy." New York: Alan R Liss, Inc., p 123.
- Nance WE (1990): Invited Editorial: Do twin Lyons have larger spots? *Am J Hum Genet* 46:646-648.
- Neilson JP, Hood VD, Cupples W, Gibson AAM, Ferguson-Smith MA (1982): Detection by ultrasound of abnormalities in twin pregnancies during the second trimester. *Br J Obstet Gynaecol* 89:1035-1040.
- Newman B, Selby JV, Quesenberry GP, King M-C, Friedman GD, Fabsitz RR (1990): Nongenetic influences of obesity on other cardiovascular disease risk factors. An analysis of identical twins. *Am J Public Health* 80:675-678.
- Northrup H, Wheless JW, Leins RA, Bertin TK (1990): Monozygotic twins discordant for tuberous sclerosis. *Am J Hum Genet* 47:A69.
- Patten RM, Mack LA, Nyberg DA, Filly RA (1989): Twin embolization syndrome: Prenatal sonographic detection and significance. *Radiology* 173:685-689.
- Pedersen IK, Philip J, Sele V, Starup J (1980): Monozygotic twins with dissimilar phenotypes and chromosome complement. *Acta Obstet Gynecol Scand* 59:459-462.
- Perlman E, Stetten G, Tuck-Muller CM, Farber RA, Neuman WL, Blakemore KJ, Hutchins GM (1990): Sexual discordance in monozygotic twins. *Am J Med Genet* 37:551-557.
- Phillips DIW (1993): Twin studies in medical research. Can they tell us whether diseases are genetically determined? *Lancet* 341:1008-1009.
- Pysher T (1980): Discordant malformations in monozygous twins. The amniotic band disruption complex. *Diagn Gynecol Obstet* 2:221-225.
- Reiss AL, Freund L, Plotnick L, Baumgardner T, Green K, Sozer AC, Reader M, Boehm C, Denckla MB (1993): The effects of X-monosomy on brain development. *Ann Neurol* 34:95-107.
- Reindollar RH, Byrd JR, Hahn DH, Haseltine FP, McDonough PG (1987): A cytogenetic and endocrinologic study of a set of monozygotic isokaryotic 45,X/46,XY twins discordant for sex. Mosaicism versus chimerism. *Fertil Steril* 47:626-633.
- Reveley AM, Reveley MA, Clifford CA, Murray RM (1982): Cerebral ventricular size in twins discordant for schizophrenia. *Lancet* i:540-541.
- Richards CS, Watkins SC, Hoffman EP, Schneider NR, Milsark IW, Katz KS, Cook JD, Kunkel LM, Cortada JM (1990): Skewed X inactivation in a female MZ twin results in Duchenne muscular dystrophy. *Am J Hum Genet* 46:672-681.
- Ross GT, Tjio JH, Lipsett MB (1969): Cytogenetic studies of presumptively monozygotic twin girls discordant for gonadal dysgenesis. *J Clin Endocrinol* 29:440-445.
- Rogers JG, Voullaire L, Gold H (1982): Monozygotic twins discordant for trisomy 21. *Am J Med Genet* 11:143-146.
- Schinzl AAGL, Smith DW, Miller JR (1979): Monozygotic twinning and structural defects. *J Pediatr* 95:921-930.
- Schlessel JS, Brown WT, Lysikiewicz A, Schiff R, Zaslav AL (1990): Monozygotic twins with trisomy 18. A report of discordant phenotype. *J Med Genet* 27:640-642.
- Schmidt R, Sobel EH, Nitowsky HM, Dar H, Allen FH (1976): Monozygotic twins discordant for sex. *J Med Genet* 13:64-67.
- Segerberg EC, Grubb WG, Henderson AE (1992): The first successful parathyroid transplant from an identical twin for the cure of permanent postoperative hypoparathyroidism. *Surgery* 111:357-358.
- Seo JW, Shin SS, Chi JG (1985): Cardiovascular system in conjoined twins: Analysis of 14 Korean cases. *Teratology* 32:151-161.
- Serville F, Carles D, Maroteaux P (1984) (Letter): Thanatophoric dysplasia in identical twins. *Am J Med Genet* 17:703-706.
- Shapiro LR, Farnsworth RG (1972): Down's syndrome in twins. *Clin Genet* 3:364-370.
- Silber SJ (1978): Transplantation of a human testis for anorchia. *Fertil Steril* 30:181-187.
- Siplovich L, Caroni R, Bar-Ziv J, Karplus M, Mares AJ (1983): Discordant Hirschsprung's disease in monozygotic twins. *J Pediatr Surg* 18:639-640.
- Sperber GH, Machin GA (1987): Microscopic study of midline determinants in janiceps twins. *March of Dimes: Birth Defects, Original Article Series* 23(1):243-275.
- Stocker JT, Heifetz SA (1987): Sirenomelia: A morphological study of 33 cases and review of the literature. *Perspect Pediatr Pathol* 10:7-50.
- Stirratt GM, Turnbull AC, Bennett J, Bobrow M, Lindenbaum RH, Wald NJ, Cuckle HS (1979): Clinical dilemmas arising from the antenatal diagnosis of neural tube defects. *Br J Obstet Gynaecol* 86:161-166.
- Teeuw AH, Kok JH (1992): Complete situs inversus in one half of a monozygotic twin pair. *Tijdschr Kindergeneesk* 60:67-69.
- Tilney NL (1986): Renal transplantation between identical twins: A review. *World J Surg* 10:381-388.
- Torgersen J (1950): Situs inversus, asymmetry, and twinning. *Am J Hum Genet* 2:361-370.

- Trejo V, Derom C, Vlietinck R, Ollier W, et al. (1994): X-chromosome inactivation patterns correspond with fetal-placental anatomy in monozygotic twin pairs: Implications for immune relatedness and concordance for autoimmunity. *Molec Med* 1:62-70.
- Tsukamoto H, Inui K, Taniike M, Kamiyama K, Hori M, Sumi K, Okida S (1993): Different clinical features in monozygotic twins. A case of 7q2 syndrome. *Clin Genet* 63:139-142.
- Tuckerman E, Webb T, Bunday SE (1985): Frequency and replication status of the fragile X fra(X)(q27-28), in a pair of monozygotic twins of a markedly differing intelligence. *J Med Genet* 22:85-91.
- Tuerlings JHAM, Breed ASPM, Vosters R, Anders GJPA (1993): Evidence of a second gamete fusion after the first cleavage of the zygote in a 47,XX,1 18/70,XXX,1 18 mosaic. A remarkable diploid-triploid discrepancy after CVS. *Prenat Diagn* 13:301-306.
- Turpin R, Lejeune J, Lafoucade J, Chigot PL, Salmon C (1961): Presumption de monozygotisme en depit d'un dimorphisme sexuel: Sujet masculin XY et sujet neutre haplo X. *Compte Rend Acad Sci* 252:2945-2946.
- Uchica IA, Freeman VCP, Gedeon M, Goldmaker J (1983): Twinning rate in spontaneous abortions. *Am J Hum Genet* 35:987-993.
- Van Itallie TB, Stunkard AJ (1990): Editorial. Using nature to understand nurture. *Am J Public Health* 80:657-658.
- Vaughn AJ, Bachman D, Sommer A (1981): Neurofibromatosis in twins: A case report of spontaneous mutation. *Am J Med Genet* 8:155-158.
- Van Thienen MN, Van der Auwerea BJ (1994): Monozygotic twins discordant for spondylocostal dysostosis. *Am J Med Genet* 52:483-486.
- Wadeluis C, Lindstedt M, Pigg M, Egberg N, Patterson U, Anvret M (1993): Hemophilia B in a 46,XX female probably caused by non-random X inactivation. *Clin Genet* 43:1-4.
- Wagner DS, Klein RL, Robinson MP, Novak RW (1990): Placental emboli from a fetus papyraceus. *J Pediatr Surg* 25:538-542.
- Walker NF (1950): Discordant monozygotic twins with retinoblastoma. *Am J Hum Genet* 2:375-384.
- Wapner RJ, Davis GH, Johnson A, Weinblatt VJ, Fischer RL, Jackson LG, Chervenak FA, McCullough LB (1990): Selective reduction of multifetal pregnancies. *Lancet* 335:90-93.
- Watson WJ, Katz VL, Albright SG, Rao KW, Aylsworth AS (1990): Monozygotic twins discordant for partial trisomy 1. *Obstet Gynecol* 76:949-951.
- Weinstrom KD, Tessen JA, Zlatnik FJ, Sipes SL (1992): Frequency, distribution, and theoretical mechanisms of hematologic and weight discordance in monochorionic twins. *Obstet Gynecol* 80:257-261.
- Westerveld AW, Sauer EW, Klassen HJ (1986): Successful treatment of a severely burned elderly patient with homografts from her identical twin sister. *Br J Plast Surg* 39:136-138.
- Winchester B, Young E, Geddes S, Genet S, Hurst J, Middleton-Price H, Williams N, Webb M, Habel A, Malcolm S (1992): Female twin with Hunger disease due to non-random inactivation of the X-chromosome: A consequence of twinning. *Am J Med Genet* 44:834-838.
- Winsor EJT, Brown BS, Luther ER, Heifetz SA, Welch JP (1987): Deceased co-twin as a cause of false positive amniotic fluid AFP and AChE. *Prenat Diagn* 7:485-489.
- Wolf HK, MacDonald J, Bradford WB, Lanman JT (1991): Acardius anceps with evidence of intrauterine vascular occlusion. *Pediatr Pathol* 11:143-152.
- Worthen NJ, Lawrence D, Bustillo M (1980): Amniotic band syndrome: Antepartum ultrasonic diagnosis of discordant anencephaly. *J Clin Ultrasound* 8:453-454A.
- Wulfsberg EA, Wassel WC, Polo CA (1991): Monozygotic twin girls with diploid-triploid chromosome mosaicism and cutaneous pigmentary dysplasia. *Clin Genet* 39:370-375.
- Yoshida K, Soma H (1986): Outcome of surviving co-twin of a fetus papyraceus or a dead twin fetus. *Acta Genet Med Gemellol* 35:91-98.
- Young ID, Patel I, Lamont AC (1989): Thanatophoric dysplasia in identical twins. *J Med Genet* 26:276-279.
- Yovich JL, Stanger JD, Granang A, Barter RA, Lunay G, Dawkins RL, Mulcahy MT (1984): Monozygotic twins from in vitro fertilization. *Fertil Steril* 41:833-837.
- Zellweger H (1968): Familial aggregates of the 21-trisomy syndrome. *Ann N Y Acad Sci* 155:785-792.
- Zneimer SM, Schneider NR, Richards CS (1993): In situ hybridization shows direct evidence of skewed inactivation in one of monozygotic twin females manifesting Duchenne muscular dystrophy. *Am J Med Genet* 45:601-605.